

TUSNÁDY'S INEQUALITY REVISITED

ANDREW CARTER AND DAVID POLLARD

ABSTRACT. Tusnády's inequality is the key ingredient in the KMT/Hungarian coupling of the empirical distribution function with a Brownian Bridge. We present an elementary proof of a result that sharpens the Tusnády inequality, modulo constants. Our method uses the beta integral representation of Binomial tails, simple Taylor expansion, and some novel bounds for the ratios of normal tail probabilities.

1. INTRODUCTION

In one of the most important probability papers of the last forty years, Komlós, Major, and Tusnády (1975) sketched a proof for a very tight coupling of the standardized empirical distribution function with a Brownian Bridge, a result now often referred to as the KMT, or Hungarian, construction. Their coupling greatly simplifies the derivation of many classical statistical results—see Shorack and Wellner (1986, Chapter 12 et seq.), for example.

The construction has taken on added significance for statistics with its use by Nussbaum (1996) in establishing asymptotic equivalence of density estimation and white noise models. Brown, Carter, Low, and Zhang (2003) have somewhat simplified and expanded Nussbaum's argument using our Theorem 2, via inequality (5).

At the heart of the KMT method (with refinements as in the exposition by Csörgő and Révész 1981, Section 4.4) lies the quantile coupling of the $\text{Bin}(n, 1/2)$ and $N(n/2, n/4)$ distributions, which may be defined as follows. Let Y be a random variable distributed $N(n/2, n/4)$. Find the cutpoints $-\infty = \beta_0 < \beta_1 < \dots < \beta_n < \beta_{n+1} = \infty$ for which

$$\mathbb{P}\{\text{Bin}(n, 1/2) \geq k\} = \mathbb{P}\{Y > \beta_k\} \quad \text{for } k = 0, 1, \dots, n.$$

When $\beta_k < Y \leq \beta_{k+1}$, let X take the value k . Then X has a $\text{Bin}(n, 1/2)$ distribution.

It is often more convenient to work with the the tails of the standard normal $\bar{\Phi}(z) = \mathbb{P}\{N(0, 1) > z\}$, and the standardized cutpoint $z_k = 2(\beta_k - n/2)/\sqrt{n}$, thereby replacing $\mathbb{P}\{Y > \beta_k\}$ by $\bar{\Phi}(z_k)$.

Symmetry considerations show that $\beta_{n-k+1} = n - \beta_k$, so that it suffices to consider only half the range for k . More precisely, when n is even, say $n = 2m$, the interval (β_m, β_{m+1}) is symmetric about $n/2$, so we have only to consider $k \geq m + 1 = (n + 2)/2$. When n is odd, say $n = 2m + 1$, the interval (β_m, β_{m+2}) is symmetric about $n/2 = \beta_{m+1}$, so we have only to consider $k \geq m + 2 = (n + 3)/2$.

Date: Original version July 2000; revised August 2003.

1991 Mathematics Subject Classification. Primary 62E17. Secondary: 62B15.

Key words and phrases. Quantile coupling; KMT/Hungarian construction; Tusnády's inequality; beta integral representation of Binomial tails; ratios of normal tails; equivalent normal deviate.

The usual normal approximation with continuity correction suggests that $\beta_k \approx k - 1/2$, which, if true, would bound $|X - Y|$ by a constant that does not change with n . Of course, such an approximation for all k is too good to be true, but results almost as good have been established. The most elegant version appeared in the unpublished dissertation (in Hungarian) of Tusnady (1977), whose key inequality may be expressed as the assertion

$$(1) \quad k - 1 \leq \beta_k \leq \frac{3n}{2} - \sqrt{2n(n-k)} \quad \text{for } n/2 \leq k \leq n.$$

As explained by Csorgo and Revesz (1981, Section 4.4), Tusnady’s inequality implies that $|X - n/2| \leq |Y - n/2| + 1$ and $|X - Y| \leq 1 + Z^2/8$, where Z denotes the standardized variable $(2Y - n)/\sqrt{n}$. They also noted that Tusnady’s proof of inequality (1) was “elementary”, but “not at all simple”. Bretagnolle and Massart (1989, Appendix) published another proof of Tusnady’s inequality—an exquisitely delicate exercise in elementary calculus and careful handling of Stirling’s formula to approximate individual Binomial probabilities. With no criticism intended, we note that their proof is quite difficult. More recently, Dudley (2000, Chapter 1) and Massart (2003) have reworked and refined the Bretagnolle/Massart calculations. Clearly there is a continuing perceived need for an accessible treatment of the coupling result that underlies the KMT construction.

With this paper we offer another approach, which actually leads to an improvement (modulo constants) of the Tusnady inequality. In fact, the Tusnady upper bound greatly overestimates β_k for moderate to large k . (See below.) Our method differs from that of Bretagnolle and Massart, in that we work directly with the whole tail probability. Our method is closer to that of Peizer and Pratt (1968), who suggested a Cornish-Fisher expansion of the Binomial percentiles—but, as noted by Pratt (1968, Sections 5 and 8), a rigorous proof by this method is difficult. To avoid the difficulty, Molenaar (1970, Section III.2) made a more direct calculation starting from the representation of the Binomial tail as a beta integral,

$$(2) \quad \mathbb{P}\{\text{Bin}(n, 1/2) \geq k\} = \frac{n!}{(k-1)!(n-k)!} \int_0^{1/2} t^{k-1}(1-t)^{n-k} dt.$$

He indicated that his expansion would be valid provided $|k - n/2| = O(\sqrt{n})$. Pratt seemed to be claiming validity for his expansion for the range $|k - n/2| = o(n)$, but we believe extra work is needed for $|k - n/2|$ large.

We should point out that Peizer and Pratt, and Molenaar, were actually concerned with normal approximations to distributions more general than the $\text{Bin}(n, 1/2)$ case needed for the KMT construction. We have specialized their results to this case.

Our method also starts from the integral representation (2), to derive an approximation via Laplace’s method for integrals (de Bruijn 1981, Section 4.3) using only Taylor’s theorem and Stirling’s formula (Feller 1968, Section II.9),

$$(3) \quad n! = \sqrt{2\pi} \exp\left(\left(n + \frac{1}{2}\right) \log n - n + \lambda_n\right) \quad \text{with } (12n+1)^{-1} \leq \lambda_n \leq (12n)^{-1}.$$

In fact (Komlos, Major, and Tusnady 1975, page 130), the KMT construction only needs a result like the Tusnady inequality for values of k in a range where $|2k - n| \leq \epsilon_0 n$ for some fixed $\epsilon_0 < 1$. For that range, a suitable bound can be derived from classical large deviation approximations for Binomial tails. For example, in an expanded version of the argument sketched in the 1975 paper, Major (2000) used

the large deviation approximation

$$\mathbb{P}\{X \geq k\} = \bar{\Phi}(\epsilon\sqrt{n}) \exp(A_n(\epsilon)) \quad \text{where } \epsilon = (2k - n)/n,$$

with

$$|A_n(\epsilon)| = O\left(n\epsilon^3 + n^{-1/2}\right) \quad \text{uniformly in } 0 \leq \epsilon \leq \epsilon_0 < 1.$$

Mason (2001) derived the KMT coupling from an analogous approximation with

$$A_n(\epsilon) = n\epsilon^3\lambda(\epsilon) + O\left(\epsilon + n^{-1/2}\right) \quad \text{uniformly in } 0 \leq \epsilon \leq \epsilon_0 < 1,$$

where $\lambda(\cdot)$ is a power series whose coefficients depend on the cumulants of the Binomial distribution. Such an approximation follows from a minor variation on the general method explained by Petrov (1975, Section 8.2). Symmetry of the $\text{Bin}(n, 1/2)$ makes the third cumulant zero; the power series $\epsilon^3\lambda(\epsilon)$ starts with a multiple of ϵ^4 .

Our method gives a sharper approximation to the $\text{Bin}(n, 1/2)$ tails over the range $n/2 < k \leq n - 1$ (which, by symmetry, actually covers the range $0 < k < n$). Only at the extreme, $k = n$, does the calculation fail.

Theorem 1. *Let X have a $\text{Bin}(n, 1/2)$ distribution, with $n \geq 28$. Define*

$$\gamma(\epsilon) = \frac{(1 + \epsilon) \log(1 + \epsilon) + (1 - \epsilon) \log(1 - \epsilon) - \epsilon^2}{2\epsilon^4} = \sum_{r=0}^{\infty} \epsilon^{2r} / (2r + 3)(2r + 4),$$

an increasing function with $\gamma(0) = 1/12$ and $\gamma(1) = -1/2 + \log 2 \approx 0.1931$. Define $\epsilon = (2K - N)/N$ where $K = k - 1$ and $N = n - 1$. Define λ_n as in (3). Then there is a constant C such that

$$\mathbb{P}\{X \geq k\} = \bar{\Phi}(\epsilon\sqrt{N}) \exp(A_n(\epsilon))$$

where

$$A_n(\epsilon) = -N\epsilon^4\gamma(\epsilon) - \frac{1}{2} \log(1 - \epsilon^2) - \lambda_{n-k} + r_k \quad \text{and} \quad -C \log N \leq Nr_k \leq C$$

for all ϵ corresponding to the range $n/2 < k \leq n - 1$.

Notice that the λ_{n-k} can be absorbed into the error terms, and that $\log(1 - \epsilon^2)$ is small compared with $N\epsilon^4 + O(n^{-1})$, when $\epsilon \leq \epsilon_0 < 1$.

A very precise approximation for the cutpoints β_k follows from inequalities (see Section 3) for the tails of the normal distribution Theorem 1.

Theorem 2. *Let $z_k = 2(\beta_k - n/2)/\sqrt{n}$ and $\epsilon = (2K - N)/N$. Let $S(\epsilon) = \sqrt{1 + 2\epsilon^2\gamma(\epsilon)}$ for $\gamma(\epsilon)$ as in Theorem 1. Then, for some constant C' and $n \geq 28$,*

$$z_k = \epsilon\sqrt{N}S(\epsilon) + \frac{\log(1 - \epsilon^2) + 2\lambda_{n-k}}{2\epsilon\sqrt{N}S(\epsilon)} + \theta_k$$

with $-C' \left(\epsilon\sqrt{N} + 1\right) \leq N\theta_k \leq C' \left(\epsilon\sqrt{N} + \log N\right)$ for all ϵ corresponding to the range $n/2 < k \leq n - 1$.

For example, the Theorem implies $\beta_k - k + 1/2 = o(1)$ uniformly over a range where $|k - n/2| = o(n^{2/3})$. Also, when $\epsilon \leq \epsilon_0 < 1/2$ the log term can be absorbed into the $O(\epsilon/\sqrt{n})$ errors. Even when k gets close to $n - 1$, the log term contributes

only a $O(n^{-1/2} \log n)$ to the approximation. More precisely, if $k = n - B$ for a fixed $B \geq 1$, our approximation simplifies to

$$(4) \quad \beta_{n-B} = \frac{1+c}{2}n - \frac{1+2B}{4c} \log n + O(1) \quad \text{where } c = S(1) \approx 1.177,$$

which agrees up to $O(1)$ terms with the result obtained by direct calculation from

$$\mathbb{P}\{X \geq n - B\} = \left(\binom{n}{0} + \cdots + \binom{n}{B} \right) 2^{-n} = \frac{n^B}{B!2^n} (1 + o(1))$$

and the well known approximation for normal percentiles,

$$\bar{\Phi}^{-1}(p) = y - \frac{\log y}{y} + O(1/y) \quad \text{as } p \rightarrow 0, \text{ where } y = \sqrt{2 \log(1/p)}.$$

By contrast, the upper bound for β_{n-B} from (1) is about $0.088n$ too large.

It is also an easy consequence of Theorem 2 that there exist positive constants C_i for which

$$(5) \quad -\frac{C_1}{\sqrt{n}} + C_2 \frac{|k - n/2|^3}{n^2} \leq \beta_k - k + \frac{1}{2} \leq \frac{C_3 \log n}{\sqrt{n}} + C_4 \frac{|k - n/2|^3}{n^2}$$

for $n/2 \leq k \leq n$ and all n . For the quantile coupling between an X distributed $\text{Bin}(n, 1/2)$ and a $Y = n/2 + \sqrt{n}Z/2$ distributed $N(n/2, n/4)$, it follows that there is a positive constant C for which

$$\left| X - \frac{n}{2} \right| \leq C + \left| Y - \frac{n}{2} \right| \quad \text{and} \quad |X - Y| \leq C + \frac{C}{n^2} \left| X - \frac{n}{2} \right|^3.$$

Using the fact that $|X - n/2| \leq n/2$, we could also write the upper bound for $|X - Y|$ as a constant multiple of $1 + Z^2 (1 \wedge |Z|/\sqrt{n})$, which improves on Tusnády's $1 + Z^2/8$, modulo multiplicative constants. (We have made no attempt to find the best constants, even though, in principle, explicit values could be found by our method.)

2. OUTLINE OF OUR METHOD

As in Theorem 1, write $\epsilon = (2K - N)/N$ where $K = k - 1$ and $N = n - 1$. Then $K/N = (1 + \epsilon)/2$ and the range $n/2 < k < n$ corresponds to

$$(6) \quad 1 - \frac{2}{N} \geq \epsilon = \frac{2K}{N} - 1 \geq \begin{cases} N^{-1} & \text{when } n \text{ is even} \\ 2N^{-1} & \text{when } n \text{ is odd.} \end{cases}$$

Define $2H(t) = (1 + \epsilon) \log t + (1 - \epsilon) \log(1 - t)$ for $0 < t < 1$. Representation (2) can then be rewritten as

$$\begin{aligned} \mathbb{P}\{X \geq k\} &= \frac{nN!}{K!(N-K)!} \int_0^{1/2} \exp(K \log t + (N-K) \log(1-t)) dt \\ &= \frac{nN!}{K!(N-K)!} \int_0^{1/2} e^{NH(t)} dt. \end{aligned}$$

By Stirling's formula (3),

$$\frac{N!}{K!(N-K)!} = \frac{1}{N} \sqrt{\frac{4N}{2\pi(1-\epsilon^2)}} \exp(\Lambda - NH(K/N)) \quad \text{where } \Lambda := \lambda_N - \lambda_K - \lambda_{N-K}.$$

Thus the beta integral equals

$$\frac{n}{N} \exp\left(\Lambda - \frac{1}{2} \log(1 - \epsilon^2) - NH(K/N)\right) \sqrt{\frac{4N}{2\pi}} \int_0^{1/2} e^{NH(t)} dt.$$

The function $H(\cdot)$ is concave on $(0, 1)$. It achieves its global maximum at K/N , which lies outside the range of integration. On the interval $(0, 1/2]$ the maximum is achieved at $1/2$. On the range of integration, $H(t) - H(K/N)$ is never greater than

$$H(1/2) - H(K/N) = -\frac{1}{2}(1 + \epsilon) \log(1 + \epsilon) - \frac{1}{2}(1 - \epsilon) \log(1 - \epsilon) = -\frac{1}{2}\epsilon^2 - \epsilon^4\gamma(\epsilon).$$

The concave function $h(s) := H((1 - s)/2) - H(1/2)$ achieves its maximum value of zero at $s = 0$ and

$$(7) \quad \mathbb{P}\{X \geq k\} = e^\Delta \sqrt{\frac{N}{2\pi}} \int_0^1 e^{Nh(s) - N\epsilon^2/2} ds$$

where $\Delta = \log(1 + N^{-1}) + \Lambda - \frac{1}{2} \log(1 - \epsilon^2) - N\epsilon^4\gamma(\epsilon)$.

The Δ contributes $O(1/n) - \lambda_{n-k} - \frac{1}{2} \log(1 - \epsilon^2) - N\epsilon^4\gamma(\epsilon)$ to the $A_n(\epsilon)$ from Theorem 1. Taylor expansion of $h(s)$ about $s = 0$ and concavity of $h(\cdot)$ show that the exponent $Nh(s)$ drops off rapidly as s moves away from zero. Indeed,

$$(8) \quad \begin{aligned} h(s) &= -\epsilon s - \frac{1}{2}s^2 + \frac{1}{6}s^3 h'''(s^*) \quad \text{with } 0 < s^* < s \\ &\approx \frac{1}{2}\epsilon^2 - \frac{1}{2}(s + \epsilon)^2 \quad \text{for } s \text{ near zero.} \end{aligned}$$

See Section 4 for the more precise statement of the approximation.

Most of the contribution to the integral (7) comes from s in a small neighborhood of 0. Ignoring tail contributions to the integral, we will then have

$$(9) \quad \mathbb{P}\{X \geq k\} \approx e^\Delta \sqrt{\frac{N}{2\pi}} \int_0^\infty \exp\left(-\frac{1}{2}N(s + \epsilon)^2\right) ds = e^\Delta \bar{\Phi}\left(\epsilon\sqrt{N}\right),$$

as asserted by Theorem 1.

To derive Theorem 2 we perturb the argument $\epsilon\sqrt{N}$ slightly to absorb the factor $\exp(A_n(\epsilon))$. We seek a y for which

$$\bar{\Phi}(\epsilon\sqrt{N} + y) \approx \exp(A_n(\epsilon)) \bar{\Phi}(\epsilon\sqrt{N}) = \bar{\Phi}(z_k).$$

That is, we need

$$\bar{\Phi}(\epsilon\sqrt{N} + y)/\bar{\Phi}(\epsilon\sqrt{N}) \approx \exp\left(-N\epsilon^4\gamma(\epsilon) - \frac{1}{2} \log(1 - \epsilon^2)\right).$$

As shown in the next Section, the ratio of normal tail probabilities $\bar{\Phi}(x + y)/\bar{\Phi}(x)$ behaves like $\exp(-xy - y^2/2)$, at least when x is large. Ignore the logarithmic term for the moment. Then the heuristic suggests that we choose y to make $\epsilon\sqrt{N}y + y^2/2 \approx N\epsilon^4\gamma(\epsilon)$, that is,

$$y \approx -\epsilon\sqrt{N} + \sqrt{N\epsilon^2 + 2N\epsilon^4\gamma(\epsilon)} \quad \text{and hence} \quad z_k \approx \epsilon\sqrt{N} + y \approx \epsilon\sqrt{N} \sqrt{1 + 2\epsilon^2\gamma(\epsilon)}$$

For the rigorous proof of Theorem 2 we need to replace these heuristic approximations by inequalities giving upper and lower bounds for $\bar{\Phi}(z_k)$, then invoke the inequalities for normal tails derived in the next Section.

3. TAILS OF THE NORMAL DISTRIBUTIONS

The classical tail bounds for the normal distribution (cf. Feller 1968, Section VII.1 and Problem 7.1) show that $\bar{\Phi}(x)$ behaves roughly like the density $\phi(x)$:

$$(10) \quad \left(\frac{1}{x} - \frac{1}{x^3}\right) \phi(x) < \bar{\Phi}(x) < \frac{1}{x} \phi(x) \quad \text{for } x > 0$$

$$\bar{\Phi}(x) < \frac{1}{2} \exp(-x^2/2)$$

The first upper bound is good for large x , the second for $x \approx 0$. For the proofs of both Theorem 1 (in Section 4) and Theorem 2 (in Section 5) we will need to bound the ratio $\bar{\Phi}(x+y)/\bar{\Phi}(x)$. It is possible to derive suitable bounds directly from (10), but we have found it easier to work with inequalities that interpolate smoothly between the different cases in (10). We express our results in logarithmic form, using the function $\Psi(x) := -\log \bar{\Phi}(x)$ and its derivative

$$\rho(x) = \frac{d}{dx} \Psi(x) = \phi(x)/\bar{\Phi}(x).$$

To a first approximation, the positive function $\rho(x)$ increases like x . By inequality (10), the error of approximation, $r(x) := \rho(x) - x$, is positive for $x > 0$ and, for $x > 1$,

$$r(x) < \frac{x}{x^2 - 1} = O(1/x) \quad \text{as } x \rightarrow \infty.$$

In fact, as shown by the proof of the next lemma, $\rho(\cdot)$ is increasing and $r(\cdot)$ is decreasing and positive, on the whole real line.

Lemma 1. *The function $\rho(\cdot)$ is increasing and the function $r(\cdot)$ is decreasing, with $r(\infty) = \rho(-\infty) = 0$ and $r(0) = \rho(0) = 2/\sqrt{2\pi} \approx .7979$. For all $x \in \mathbb{R}$ and $\delta \geq 0$, the increments of the function $\Psi(x) := -\log \bar{\Phi}(x)$ satisfy the inequalities*

- (i) $\delta\rho(x) \leq \Psi(x+\delta) - \Psi(x) \leq \delta\rho(x+\delta)$,
- (ii) $\delta r(x+\delta) \leq \Psi(x+\delta) - \Psi(x) - \frac{1}{2}(x+\delta)^2 + \frac{1}{2}x^2 \leq \delta r(x)$,
- (iii) $x\delta + \frac{1}{2}\delta^2 \leq \Psi(x+\delta) - \Psi(x) \leq \rho(x)\delta + \frac{1}{2}\delta^2$.

Proof. Let Z be $N(0, 1)$ distributed. Define $M(x) = \mathbb{P}e^{-x|Z|}$, a decreasing function of x with $\log M(x)$ strictly convex. Notice that

$$1/\rho(x) = \sqrt{2\pi} \exp(x^2/2) \int_0^\infty \phi(z+x) dz = \int_0^\infty \exp(-xz - z^2/2) dz = \sqrt{\frac{\pi}{2}} M(x).$$

Thus $-\log M(x) - \log \sqrt{\pi/2} = \log \rho(x) = \Psi(x) - x^2/2 - \log \sqrt{2\pi}$ is a concave, increasing function of x with derivative $\rho(x) - x = r(x)$. It follows that $r(\cdot)$ is a decreasing function, because

$$r'(x) = -\frac{d^2}{dx^2} \log M(x) < 0 \quad \text{by convexity of } \log M(x).$$

Inequality (i) follows from the equality

$$\Psi(x+\delta) - \Psi(x) = \delta\Psi'(y^*) = \delta\rho(y^*) \quad \text{for some } x < y^* < x+\delta,$$

together with the fact that $\rho(\cdot)$ is an increasing function. Similarly, the fact that

$$\frac{d}{dy} (\Psi(y) - \frac{1}{2}y^2) = \rho(y) - y = r(y) \quad \text{which is a decreasing function}$$

gives inequality (ii). Inequality (iii) follows from (ii) because $\delta r(x+\delta) \geq 0$ and $x\delta + r(x)\delta = \rho(x)\delta$. \square

Reexpressed in terms of the tail function $\bar{\Phi}$, the three inequalities from the lemma become

- (i) $\exp(-\delta\rho(x)) \geq \bar{\Phi}(x+\delta)/\bar{\Phi}(x) \geq \exp(-\delta\rho(x+\delta))$,
- (ii) $\exp(-\delta r(x+\delta)) \geq \exp(x\delta + \delta^2/2) \bar{\Phi}(x+\delta)/\bar{\Phi}(x) \geq \exp(-\delta r(x))$,
- (iii) $\exp(-x\delta - \delta^2/2) \geq \bar{\Phi}(x+\delta)/\bar{\Phi}(x) \geq \exp(-\rho(x)\delta - \delta^2/2)$.

Less formally,

$$\mathbb{P}\{Z \leq x + \delta \mid Z \leq x\} = 1 - \bar{\Phi}(x + \delta)/\bar{\Phi}(x) \approx \delta\rho(x) \quad \text{for small } \delta,$$

which corresponds to the fact that ρ is the hazard rate for the $N(0, 1)$ distribution.

4. DETAILS OF THE PROOF FOR THEOREM 1

To make the proof rigorous, we need to replace the approximation in the Taylor expansion (8) by upper and lower bounds involving the third derivative

$$h'''(s) = \frac{1 - \epsilon}{(1 + s)^3} - \frac{1 + \epsilon}{(1 - s)^3} = -\frac{6s + 2s^2 + \epsilon(2 + 6s^2)}{(1 - s^2)^3}.$$

The derivative of this function is negative for all s . Thus

$$h'''(s) \leq h'''(0) = -2\epsilon \quad \text{for } 0 < s < 1$$

and

$$h(s) \leq \frac{1}{2}\epsilon^2 - \frac{1}{2}(s + \epsilon)^2 \quad \text{for } 0 < s < 1.$$

The right-hand side of the approximation (9) is actually an upper bound, because the integrand is nonnegative on $(1, \infty)$. That is,

$$\mathbb{P}\{X \geq k\} \leq e^{\Delta} \bar{\Phi}(\epsilon\sqrt{N}),$$

which gives the upper bound for $A_n(\epsilon)$ stated in the Theorem.

For the lower bound, for some small positive η discard the contribution to the integral in (7) from the range $(\eta, 1)$, and bound h''' from below by $h'''(\eta)$ on the range $(0, \eta)$, then integrate to get

$$\begin{aligned} \mathbb{P}\{X \geq k\} &\geq e^{\Delta} \sqrt{\frac{N}{2\pi}} \int_0^{\eta} \exp\left(-\frac{1}{2}N(s + \epsilon)^2 + \frac{1}{6}N\eta s^2 h'''(\eta)\right) ds \\ &= e^{\Delta} \sqrt{\frac{N}{2\pi}} \int_0^{\eta} \exp\left(-\frac{1}{2}N\kappa^2(s + \epsilon/\kappa^2)^2 + \frac{1}{2}N\epsilon^2/\kappa^2 - \frac{1}{2}N\epsilon^2\right) ds \\ &= \frac{e^{\Delta}}{\kappa} \exp\left(\frac{1}{2}N\epsilon^2/\kappa^2 - \frac{1}{2}N\epsilon^2\right) \left(\bar{\Phi}(\epsilon\sqrt{N}/\kappa) - \bar{\Phi}(\epsilon\sqrt{N}/\kappa + \kappa\eta\sqrt{N})\right) \end{aligned}$$

where

$$\kappa^2 = 1 - \frac{1}{3}\eta h'''(\eta) \leq 1 + 6\eta(\eta + \epsilon) \quad \text{if } \eta \leq \frac{1}{2}.$$

From Lemma 1, parts (iii) and (ii),

$$\bar{\Phi}(\epsilon\sqrt{N}/\kappa + \kappa\eta\sqrt{N}) \leq \bar{\Phi}(\epsilon\sqrt{N}/\kappa) \exp(-N\epsilon\eta - \frac{1}{2}N\kappa^2\eta^2)$$

and

$$\exp\left(\frac{1}{2}N\epsilon^2\right) \bar{\Phi}(\epsilon\sqrt{N}) \leq \exp\left(\frac{1}{2}N\epsilon^2/\kappa^2\right) \bar{\Phi}(\epsilon\sqrt{N}/\kappa).$$

Thus

$$(11) \quad \mathbb{P}\{X \geq k\} \geq \exp(\Delta - \log \kappa) \bar{\Phi}(\epsilon\sqrt{N}) \left[1 - \exp(-N\epsilon\eta - \frac{1}{2}N\kappa^2\eta^2)\right].$$

We need $\log \kappa = O(\ell_N)$, where $\ell_N = N^{-1} \log N$, for otherwise the asserted inequality $-C \log N \leq r_k$ would be violated. As $\log \kappa \leq 6(\eta^2 + \eta\epsilon)$, this requirement

suggest that we take η as a solution to the equation $\frac{1}{2}\eta^2 + \eta\epsilon = \ell_N$, that is, $\eta := -\epsilon + \sqrt{\epsilon^2 + 2\ell_N}$. We would then have $\kappa^2 \leq 1 + 12\ell_N$ and $\eta \leq 1/2$, at least for $n \geq 28$. Also, the exponent $-N\epsilon\eta - \frac{1}{2}N\kappa^2\eta^2$ is smaller than $-\log N$, which ensures that the final, bracketed term in (11) only contributes another $O(N^{-1})$ to the $A_n(\epsilon)$ from Theorem 1.

5. DETAILS OF THE PROOF FOR THEOREM 2

Written using the Ψ function from Lemma 1, the assertion of Theorem 1 implies that

$$\Psi(z_k) = \Psi(\epsilon\sqrt{N}) + B_n(\epsilon) + \tau_k$$

where, for some constant C ,

$$B_n(\epsilon) = N\epsilon^4\gamma(\epsilon) + \frac{1}{2}\log(1 - \epsilon^2) + \lambda_{n-k} \quad \text{and} \quad -CN^{-1} \leq \tau_k \leq C\ell_N$$

for ϵ corresponding to the range $n/2 \geq k \leq n - 1$. That is, for $0 \leq \epsilon \leq 1 - 2N^{-1}$.

Define

$$w_k = \epsilon\sqrt{N}S(\epsilon) + \frac{\log(1 - \epsilon^2) + 2\lambda_{n-k}}{2\epsilon\sqrt{N}S(\epsilon)}.$$

We need to show that there is a constant C' for which $z_k = w_k + \theta_k$, with $-C'(\epsilon\sqrt{N} + 1) \leq N\theta_k \leq C'(\epsilon\sqrt{N} + \log N)$ for $0 \leq \epsilon \leq 1 - 2N^{-1}$. Consider two cases.

5.1. Suppose $\epsilon \leq C_0/\sqrt{N}$, for some constant C_0 . Uniformly over that range, $B_n(\epsilon) = O(N^{-1})$ and $w_k = \epsilon\sqrt{N} + O(N^{-1})$. From Lemma 1(i), for all nonnegative δ_1 and δ_2 ,

$$\Psi(x) + \delta_1\rho(x) \leq \Psi(x + \delta_1) \quad \text{and} \quad \Psi(x - \delta_2) + \delta_2\rho(x - \delta_2) \leq \Psi(x).$$

With x equal to $\epsilon\sqrt{N}$ and C_1 a large enough constant, deduce that

$$\Psi(\epsilon\sqrt{N} - C_1N^{-1}) < \Psi(z_k) < \Psi(\epsilon\sqrt{N} + C_1\ell_N)$$

and hence

$$w_k - O(N^{-1}) - C_1N^{-1} < z_k < w_k + O(N^{-1}) + C_1\ell_N$$

5.2. Suppose $C_0/\sqrt{N} \leq \epsilon \leq 1 - 2N^{-1}$. Write x for $\epsilon\sqrt{N}$ and β for $B_n(\epsilon) + \tau_k = \Psi(z_k) - \Psi(x)$. For all ϵ in this range, if C_0 is large enough we have $\beta > 0$ and $r(x) \leq 2/x$. The function $h(t) = t - \sqrt{t^2 + 2\beta}$ is negative, increasing and concave, with $h'(t) \leq 2\beta/t^2$. The positive numbers $\delta_1 = -h(x)$ and $\delta_2 = -h(\rho(x))$ are roots of two quadratic equations, $\delta_1x + \frac{1}{2}\delta_1^2 = \beta = \delta_2\rho(x) + \frac{1}{2}\delta_2^2$. From Lemma 1(iii),

$$\begin{aligned} \Psi(z_k) - \Psi(x) &= x\delta_1 + \frac{1}{2}\delta_1^2 \leq \Psi(x + \delta_1) - \Psi(x), \\ \Psi(x + \delta_2) - \Psi(x) &\leq \rho(x)\delta_2 + \frac{1}{2}\delta_2^2 = \Psi(z_k) - \Psi(x), \end{aligned}$$

which imply that $x + \delta_2 \leq z_k \leq x + \delta_1$. These bounds force z_k to lie close to $x + \delta_1$:

$$0 \leq x + \delta_1 - z_k \leq \delta_1 - \delta_2 = h(\rho(x)) - h(x) \leq r(x)h'(x) \leq 4\beta/x^3 = O(\epsilon/\sqrt{N}).$$

And $x + \delta_1$ lies close to w_k :

$$\begin{aligned} x + \delta_1 &= \sqrt{N\epsilon^2 + 2\beta} \\ &= \epsilon\sqrt{N} S(\epsilon) \left(1 + \frac{\log(1 - \epsilon^2) + 2\lambda_{n-k} + \tau_k}{N\epsilon^2 S(\epsilon)^2} \right)^{1/2} \\ &= w_k + \frac{\tau_k}{2\epsilon\sqrt{N} S(\epsilon)} + O\left(\sqrt{N}\ell_N^2\right). \end{aligned}$$

The assertion of Theorem 2 follows.

ACKNOWLEDGMENTS.

We thank the referees for particularly constructive comments and suggested improvements, David Mason for explaining the role of large deviation approximations, and Sandor Csörgő for providing, many years ago, an English translation of parts of Tusnády's dissertation.

REFERENCES

- Bretagnolle, J. and P. Massart (1989). Hungarian constructions from the nonasymptotic viewpoint. *Annals of Probability* 17, 239–256.
- Brown, L. D., A. V. Carter, M. G. Low, and C.-H. Zhang (2003). Asymptotic equivalence theory for a Poisson process with variable intensity. (Submitted to *Annals of Statistics*).
- Csörgő, M. and P. Révész (1981). *Strong Approximations in Probability and Statistics*. New York: Academic Press.
- de Bruijn, N. G. (1981). *Asymptotic Methods in Analysis*. Dover.
- Dudley, R. M. (2000). Notes on empirical processes. Lecture notes for a course given at Aarhus University, August 1999.
- Feller, W. (1968). *An Introduction to Probability Theory and Its Applications* (third ed.), Volume 1. New York: Wiley.
- Komlós, J., P. Major, and G. Tusnády (1975). An approximation of partial sums of independent rv-s, and the sample df. I. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* 32, 111–131.
- Major, P. (2000). The approximation of the normalized empirical distribution function by a Brownian bridge. Technical report, Mathematical Institute of the Hungarian Academy of Sciences. Notes available from <http://www.renyi.hu/~major/>.
- Mason, D. M. (2001). Notes on the KMT Brownian bridge approximation to the uniform empirical process. In N. Balakrishnan, I. A. Ibragimov, and V. B. Nevzorov (Eds.), *Asymptotic Methods in Probability and Statistics with Applications*, pp. 351–369. Boston: Birkhäuser.
- Massart, P. (2003). Strong approximation for multivariate empirical and related processes, via KMT constructions. *Annales de l'Institut Henri Poincaré* 38, 991–1007.
- Molenaar, W. (1970). *Approximations to the Poisson, Binomial, and Hypergeometric distribution functions*. Center for Mathematics and Computer Science. CWI Tract 31.
- Nussbaum, M. (1996). Asymptotic equivalence of density estimation and gaussian white noise. *Annals of Statistics* 24, 2399–2430.

- Peizer, D. B. and J. W. Pratt (1968). A normal approximation for Binomial, F, and other common, related tail probabilities I. *Journal of the American Statistical Association* 63, 1416–1456.
- Petrov, V. V. (1975). *Sums of Independent Random Variables*. Springer-Verlag. (English translation from 1972 Russian edition).
- Pratt, J. W. (1968). A normal approximation for Binomial, F, and other common, related tail probabilities II. *Journal of the American Statistical Association* 63, 1457–1483.
- Shorack, G. R. and J. A. Wellner (1986). *Empirical Processes with Applications to Statistics*. New York: Wiley.
- Tusnády, G. (1977). *A study of Statistical Hypotheses*. Ph. D. thesis, Hungarian Academy of Sciences, Budapest. In Hungarian.

ANDREW CARTER, DEPARTMENT OF STATISTICS AND APPLIED PROBABILITY, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CALIFORNIA 93106.

E-mail address: carter@pstat.ucsb.edu

URL: <http://www.pstat.ucsb.edu/faculty/carter/>

DAVID POLLARD, STATISTICS DEPARTMENT, YALE UNIVERSITY, BOX 208290 YALE STATION, NEW HAVEN, CT 06520-8290.

E-mail address: david.pollard@yale.edu

URL: <http://www.stat.yale.edu/~pollard/>